

COMMON FIXED POINT THEOREM OF FOUR MAPPING

PRASHANT CHAUHAN

Department of Applied Science & Humanities, C. E. T. Moradabad, Uttar Pradesh, India

ABSTRACT

Srivastava [12], [9], Dubey & Dubey [3], Bhola & Sharma [1] Pandey & Dubey [6]; Considered three self mappings and obtained a unique common fixed point. Here we generalized the contraction used by above authors for five maps and obtained a unique fixed point.

KEYWORDS: Fixed Point Theorems, Mappings, Contraction Mappings, Continuous Mappings, Complete Metric Space

INTRODUCTION

Srivastava [9] used the inequality as follows

$$[1.1] \quad d(E_x, f_y) \leq C_1 \left[\frac{d(T_x, E_x)d(T_y, F_y)}{d(T_x, T_y)} \right] + C_2 \left[\frac{d(T_x, F_y)d(T_y, F_y)}{d(T_x, F_y)} \right] \\ + C_3 [d(T_x, F_y) + d(T_y, F_y)] + C_4 [d(T_x, F_y) + d(T_y, E_x)] \\ + C_5 [d(T_x, T_y) + d(E_x, F_y)] + C_6 d(T_x, T_y); \quad \forall x, y \in X$$

$$T_x \neq T_y$$

Then, E, F, T have a unique common fixed point if

$$C_i \geq 0; \quad 0 \leq C_1 + 2(C_2 + C_3 + C_4 + C_5 + C_6) \leq 1,$$

$$0 \leq 2(C_4 + C_5 + C_6) \leq 1;$$

Dubey & Dubey [3] have proved a fixed point theorem for three maps S, T & I of a complete metric space (X, d) satisfying:

$$[1.2] \quad d(S_x, T_y) \leq \frac{q \{ \alpha d(I_x, S_x) d(T_x, T_y) + \beta d(T_y, S_x) d(I_y, T_y) + \gamma d(I_x, T_y) \}^2}{\alpha d(T_x, T_y) + \beta d(I_y, S_x) + \gamma d(I_x, T_y)}$$

$$\forall \quad \alpha d(T_x, T_y) + \beta d(I_y, S_x) + \gamma d(I_x, T_y) \neq 0$$

$$0 \leq q < 1, \quad \alpha, \beta, \gamma \geq 0$$

Pandey & Dubey [6] obtained a unique common fixed point of three maps E, F & T satisfying.

$$[1.3] \quad d(E_x, F_y) \leq \frac{\alpha_1 d(T_x, F_y) d(T_x, T_y)}{d(T_x, T_y) + d(T_x, F_y)} + \alpha_2 [d(T_x, E_x) + d(T_y, F_y)] + \alpha_3 [d(T_x, F_y) + d(T_y, F_x)] + \alpha_4 d(T_x, T_y)$$

Here we have generalized the contraction on five self maps as follows:

T_1, T_2, T_3, T_4 & T_5 on X to itself

$$[1.4] \quad d(T_1, T_2 x, T_3, T_4 y) \leq C_1 \left[\frac{d(T_5 x, T_1 T_2 x) d(T_5 y, T_3 T_4 y)}{d(T_5 x, T_5 y)} \right] + C_2 \left[\frac{d(T_5 x, T_3 T_4 y) d(T_5 y, T_3 T_4 y)}{d(T_1 T_2 x, T_3 T_4 y)} \right] \\ + C_3 [d(T_5 x, T_1 T_2 x) + d(T_5 y, T_3 T_4 y)] + C_4 [d(T_5 x, T_3 T_4 y) + d(T_5 y, T_1 T_2 x)] \\ + C_5 [d(T_5 x, T_5 y) + d(T_1 T_2 x, T_3 T_4 y)] + C_6 d(T_5 x, T_5 y).$$

We see that under certain condition we get a unique common fixed point.

MAIN RESULTS

Let (X, d) be a complete metric space & $T_i : X \rightarrow X$; $i = 1, 2, 3, 4, 5$ be five mapping satisfying [1.4] and [2.1] as follows

$$[2.1] \quad T_2 x \neq T_5 y; \quad \alpha_i \geq 0,$$

$$T_5(T_1 T_2) = (T_1 T_2)(T_5), \quad T_5(T_3 T_4) = (T_3 T_4)(T_5)$$

$$T_1 T_2 = T_2 T_1, \quad T_3 T_4 = T_4 T_3$$

$$T_1 T_2(X) \subseteq T_5(X), \quad T_3 T_4(X) \subseteq T_5(X)$$

Then $T_i (i = 1 \text{ to } 5)$ have a unique common fixed point in X .

Proof: Let $x_0, x_1, x_2 \in X$ and $T_1 T_2(X) \subseteq T_5(X)$ such that

$$T_1 T_2(x_0) = T_5(x_1), \quad T_3 T_4(x_1) = T_5(x_2)$$

$$T_3 T_4(X) \subseteq T_5(X)$$

Since $T_1 T_2 x_{2n} = T_5 x_{2n+1}$, $T_3 T_4 x_{2n+1} = T_5 x_{2n+1}$, replace $x \rightarrow x_{2n}$ & $y \rightarrow x_{2n-1}$, then we have from [1.4]

$$[2.3] \quad d(T_5 x_{2n+1}, T_5 x_{2n}) = d(T_1 T_2 x_{2n}, T_3 T_4 x_{2n-1})$$

$$d(T_1 T_2 x_{2n}, T_3 T_4 x_{2n-1}) \leq C_1 \frac{d(T_5 x_{2n}, T_1 T_2 x_{2n}) d(T_5 x_{2n-1}, T_3 T_4 x_{2n-1})}{d(T_5 x_{2n}, T_5 x_{2n-1})} \\ + C_2 \frac{d(T_5 x_{2n}, T_3 T_4 x_{2n-1}) d(T_5 x_{2n-1}, T_3 T_4 x_{2n-1})}{d(T_1 T_2 x_{2n}, T_3 T_4 x_{2n-1})}$$

$$\begin{aligned}
& + C_3 [d(T_5 x_{2n}, T_1 T_2 x_{2n}) + d(T_5 x_{2n-1}, T_3 T_4 x_{2n-1})] \\
& + C_4 [d(T_5 x_{2n}, T_3 T_4 x_{2n-1}) + d(T_5 x_{2n-1}, T_1 T_2 x_{2n})] \\
& + C_5 [d(T_5 x_{2n}, T_5 x_{2n-1}) + d(T_1 T_2 x_{2n}, T_3 T_4 x_{2n-1})] \\
& + C_6 d(T_5 x_{2n}, T_5 x_{2n-1}).
\end{aligned}$$

Which yields from (2.2) and using Property of Metric Space.

$$d(T_5 x_{2n+1}, T_5 x_{2n}) \{1 - C_1 - C_3 - C_5\} \leq d(T_5 x_{2n-1}, T_5 x_{2n}) (C_3 + C_4 + C_5)$$

Or

$$[2.4] \quad d(T_5 x_{2n+1}, T_5 x_{2n}) \leq K d(T_5 x_{2n}, T_5 x_{2n-1})$$

Where

$$0 < K = \frac{(C_3 + C_4 + C_5)}{1 - C_1 - C_3 - C_5} < 1$$

Thus $[T_5 x_{2n}]$ is a Cauchy Sequence in complete metric space X & have a common fixed point u in X such that

$$[2.5] \quad T_5 x_{2n+1} = u = \lim_{n \rightarrow \infty} T_5 x_{2n}$$

Now if $T_1 T_2 u \neq T_5 u$ then

$$\begin{aligned}
[2.6] \quad & (T_1 T_2 u, T_5 u) \leq \lim d(T_1 T_2 u, T_3 T_4 T_5 x_{2n+1}) \\
& \leq C_1 \frac{\lim d(T_5 u, T_1 T_2 u) d(T_5 T_5 x_{2n+1}, T_3 T_4 T_5 x_{2n+1})}{d(T_5 u, T_5 T_5 x_{2n+1})} \\
& + C_2 \frac{d(T_5 u, T_3 T_4 T_5 x_{2n+1}) d(T_5 T_5 x_{2n+1}, T_3 T_4 T_5 x_{2n+1})}{d(T_1 T_2 u, T_3 T_4 T_5 x_{2n+1})} \\
& + C_3 [d(T_5 u, T_1 T_2 u) + d(T_5 T_5 x_{2n+1}, T_3 T_4 T_5 x_{2n+1})] \\
& + C_4 [d(T_5 u, T_3 T_4 T_5 x_{2n+1}) + d(T_5 T_5 x_{2n+1}, T_1 T_2 u)] \\
& + C_5 [d(T_5 u, T_5 T_5 x_{2n+1}) d(T_1 T_2 u, T_3 T_4 T_5 x_{2n+1})] + C_6 [d(T_5 u, T_5 T_5 x_{2n+1})]
\end{aligned}$$

Which Reduced to

$$d(T_1 T_2 u, T_5 u) (1 - C_5 - C_4 - C_3) \leq 0.$$

$$[2.7] \quad d(T_1 T_2 u, T_5 u) \leq 0$$

Which is a Contraction therefore

$T_1T_2(u) = T_5(u)$, Also we have $T_5(u) = T_3T_4(u)$, hence

$$[2.8] \quad T_1T_2(u) = T_5(u) = T_3T_4(u), \text{ and}$$

$$[2.9] \quad \begin{aligned} T_5(T_5u) &= T_5(T_1T_2u) = T_1T_2(T_1T_2u) = T_1T_2(T_3T_4u) = T_1T_2(T_5u) = T_5(T_3T_4u) \\ &= T_3T_4(T_5u) = T_3T_4(T_1T_2u) = T_3T_4(T_3T_4u) \end{aligned}$$

Now if $T_1T_2u \neq T_3T_4(T_1T_2u)$

$$[2.10] \quad \begin{aligned} d(T_1T_2u, T_3T_4(T_1T_2u)) &\leq C_1 \frac{d(T_5u, T_1T_2u)d(T_5T_1T_2u, T_3T_4T_3T_4u)}{d(T_5u, T_5T_3T_4u)} \\ &+ C_2 \frac{d(T_5u, T_3T_4T_3T_4u)d(T_5T_3T_4u, T_3T_4T_3T_4u)}{d(T_1T_2u, T_3T_4T_3T_4u)} \\ &+ C_3 [d(T_5u, T_1T_2u) + d(T_5T_3T_4u, T_3T_4T_3T_4u)] \\ &+ C_4 [d(T_5u, T_3T_4T_3T_4u) + d(T_5T_3T_4u, T_1T_2u)] \\ &+ C_5 [d(T_5u, T_5T_3T_4u) + d(T_1T_2u, T_3T_4T_3T_4u)] + C_6 d(T_5u, T_5T_3T_4u). \end{aligned}$$

Therefore (2.10) reduces to

$$d(T_5u, T_5T_5u)(1 - 2C_4 - 2C_5 - C_6) \leq 0$$

Or equivalently

$$(2.11) \quad d(T_5u, T_5T_5u) \leq 0$$

Which is a Contraction, therefore

$$T_5(T_5(u)) = T_5(u)$$

Also

$$T_3T_4(T_1T_2u) = T_1T_2u$$

Thus we get

Thus $T_1T_2u = v$ is a common fixed point of $T_5T_1T_2$ & T_3T_4 , if possible we take another common fixed point of $T_1T_2T_5$ & T_3T_4 . Thus

$$[d(v, w)] = [d(T_1T_2v, T_3T_4w)]$$

[2.12] which is a view of [1.4] yields

$$\begin{aligned}
d(v, w) \leq & C_1 \frac{[d(T_5 v, T_1 T_2 v)]d(T_5 w, T_5 T_4 w)}{d(T_5 v, T_5 w)} + C_2 \frac{[d(T_5 v, T_3 T_4 w)d(T_5 w, T_3 T_4 w)]}{d(T_1 T_2 v, T_3 T_4 w)} \\
& + C_3 [d(T_5 v, T_3 T_4 v) + d(T_5 w, T_3 T_2 w)] + C_4 [d(T_5 v, T_3 T_4 w) + d(T_5 w, T_1 T_2 v)] \\
& + C_5 [d(T_5 v, T_5 w) + d(T_1 T_2 v, T_3 T_4 w)] + C_6 d(T_5 v, T_5 w)
\end{aligned}$$

Equation [2.12] reduced to

$$d(v, w)(1 - 2C_4 - 2C_5 - C_6) \leq 0$$

Or equivalently

$$d(v, w) = 0 \Rightarrow v = w$$

Now to show that v is a unique common fixed point of T_1, T_2, T_3, T_4 & T_5 we find that

$$T_1(v) = T_1(T_1 T_2 v) = T_1(T_2 T_1 v) = T_1 T_2(T_1 v) \Rightarrow T_1 v$$

Which implies that $T_1 v$ is as another fixed point of T_1 & T_2 . Hence by uniqueness of fixed point of T_1 & T_2 , we obtain v a fixed point of T_1 , similarly a unique fixed point of T_2, T_3, T_4 & T_5 . Hence proved.

REFERENCES

1. Bhola P.K. & Sharma P.L. (1991): Common fixed point Theorem for Three Maps. Bull. Cal. Math. Soc. 83, 398-400.
2. Dinesh Rani Chug, Renu Chug & Ramesh Kumar (1996): A theorem on common fixed point for five mappings on metric space. The Mathematics Equation XXX, 1933-201.
3. Dubey R.P. & Dubey B, N. (1991): Common fixed point theorem for three mappings rational inequality. G Anita 42, 75-80.
4. M. Abbas, B.E. Rhoades (2009): Fixed and periodic point results in cone metric Spaces, Appl. Math. Lett. 22, 511-515.
5. Pachpatte, B.G. (1980): Some fixed point theorem for mapping in 2-metric space hung Yuan J; 8, 7-12.
6. Pandey P.K. & Dubey R.K. (1992): On common fixed point theorem of three mappings. Mathematics Student 61, 97-100.
7. Pathak H.K. & Dubey R.P. (1991): A common fixed point theorem for three commuting mappings in uniform spaces. The Mathematics Education XXV, 110-113.
8. Rawat Pratimma & Sharma P.L. (1980): Common fixed point theorem for three mapping Acta Ciencia Indica XVII, 351-354.
9. Srivastava Rakesh (1993): On common fixed point theorem of three mappings. Ganita 44, 81-95.
10. S. Rezapour, Halbarani (2008): Some notes on the paper "cone metric spaces and fixed point theorem of contractive mappings", J. Math. Anal. Appl., 345, 719-724.

11. Yian Zhao ,Guang Xing Song, X. Huang, Xiaoyan Sun, Guotao Wang, 23 (2010): New common fixed point theorems for maps on cone metric spaces, Applied Mathematics Letters, 1033-1037.
12. Srivastava Rakesh (1991): Common fixed point theorem for continuous mapping Ganita 71-74.
13. V. Berinde, (2009): A common fixed point theorem for compatible quasi contractive self Mappings in Metric Spaces, Appl. Math. Computer. 213,348-354.